

PROBING THE SCALAR SECTOR IN $e^+e^- \rightarrow f\bar{f}\phi$

M. L. STONG

Department of Physics, University of Wisconsin, Madison, WI 53706, USA

and

K. HAGIWARA

Theory Group, KEK, Tsukuba, Ibaraki 305, Japan

ABSTRACT

We study possible deviations from the Standard Model in the reaction $e^+e^- \rightarrow Z\phi$, where ϕ denotes a spinless neutral boson. We show how the Z decay angular correlation can be used to extract detailed information on the ϕ couplings, such as the parity of ϕ , radiatively induced form factor effects and possible CP violation in the scalar sector.

1. Introduction and Effective Lagrangian

The process $e^+e^- \rightarrow ZH$ is expected to be the best reaction to look for the Higgs boson of mass $\lesssim 2m_W$ at LEP II and at an early stage of the next linear e^+e^- linear colliders¹. Unlike in Higgs hunting at hadron colliders², we expect to learn details of the Higgs boson properties and interactions at e^+e^- colliders. These include the search for a deviation from the minimal one-doublet Higgs boson model and for possible radiative effects³. In fact a neutral spinless boson ϕ which is produced via $e^+e^- \rightarrow Z\phi$ may not be a Higgs boson at all, but a new type of a bound state such as a pseudo Nambu-Goldstone boson⁴ of a new strong interaction with a spontaneously broken chiral symmetry. The particle would then be a pseudoscalar rather than a scalar. We may even expect to observe a CP-violating interaction in the boson sector⁵.

In this paper, we study couplings of a spinless neutral boson ϕ , which may be a scalar, a pseudoscalar, or some mixture of the two, to the Z boson in the process $e^+e^- \rightarrow (Z \rightarrow f\bar{f})\phi$ by using the angular distributions of the final-state fermions. We consider an effective Lagrangian which contains the Standard Model couplings of fermions to the Z and γ , and study the effects of general mass-dimension-five ϕZZ and $\phi Z\gamma$ couplings which respect electromagnetic gauge invariance*:

$$\mathcal{L}_{eff} = a_Z \phi Z^\mu Z_\mu + \sum_{V=Z,\gamma} V^{\mu\nu} \left[b_V \phi Z_{\mu\nu} + c_V [(\partial_\mu \phi) Z_\nu - (\partial_\nu \phi) Z_\mu] + \tilde{b}_V \phi \tilde{Z}_{\mu\nu} \right]. \quad (1)$$

The terms a_Z , b_V , and c_V alone would correspond to a CP-even scalar ϕ , while the term $\tilde{b}_V$ alone indicates a CP-odd pseudoscalar. Interference of these two sets of terms leads to CP-violating effects.

* $V_{\mu\nu} = \partial_\mu V_\nu - \partial_\nu V_\mu$, $\tilde{V}_{\mu\nu} \equiv \frac{1}{2} \varepsilon_{\mu\nu\alpha\beta} V^{\alpha\beta}$, with the convention $\varepsilon_{0123} = +1$.

These effective interactions contribute to the process $e^+e^- \rightarrow Z\phi$. For electron helicity $\sigma = \pm 1$ (in units of $\hbar/2$) and Z polarization λ , the helicity amplitudes for $e^+e^- \rightarrow Z\phi$ are given by:

$$\begin{aligned}\mathcal{M}_\sigma^{(\lambda=0)} &= \widehat{M}_\sigma^{(\lambda=0)} \sigma \sin \theta, \\ \mathcal{M}_\sigma^{(\lambda=\pm 1)} &= \widehat{M}_\sigma^{(\lambda=\pm 1)} \frac{1 + \lambda \sigma \cos \theta}{\sqrt{2}},\end{aligned}\quad (2)$$

where

$$\begin{aligned}\widehat{M}_\sigma^{(\lambda=0)} &= \frac{g_Z (v_e + \sigma a_e)}{s - m_Z^2 + im_Z \Gamma_Z} \left(2\sqrt{s} \frac{\omega}{m_Z} (a_Z + (s + m_Z^2)c_Z) + 4sm_Z(b_Z - c_Z) \right) \\ &\quad - e \left(2m_Z(b_\gamma - c_\gamma) + 2\frac{\sqrt{s}\omega}{m_Z} c_\gamma \right),\end{aligned}\quad (3)$$

$$\begin{aligned}\widehat{M}_\sigma^{(\lambda=\pm 1)} &= \frac{g_Z (v_e + \sigma a_e)}{s - m_Z^2 + im_Z \Gamma_Z} \left(2\sqrt{s} (a_Z + (s + m_Z^2)c_Z) + 4s\omega(b_Z - c_Z) - i\lambda 4sk\tilde{b}_Z \right) \\ &\quad - e \left(2\omega(b_\gamma - c_\gamma) + 2\sqrt{s}c_\gamma - i\lambda 2k\tilde{b}_\gamma \right).\end{aligned}\quad (4)$$

Here θ is the polar angle of the Z momentum about the electron beam direction, and ω and k are the Z energy and momentum in the e^+e^- c.m. frame. The couplings are denoted by $e = \sqrt{4\pi\alpha}$, $g_Z = e/(\sin \theta_W \cos \theta_W)$, $v_e = -1/4 + \sin^2 \theta_W$, and $a_e = 1/4$. The total cross section is simply obtained from the matrix elements of Eq. 4, but does not allow us to separate contributions from the various couplings.

3. Decay Angular Correlations

The differential cross-section for a given electron (f) helicity σ (σ') is

$$\frac{d\sigma(\sigma, \sigma')}{d\cos\theta d\cos\hat{\theta} d\hat{\varphi}} = \frac{1}{32\pi s} \bar{\beta} \left(\frac{m_Z^2}{s}, \frac{m_\phi^2}{s} \right) \frac{g_Z^2 (v_f + \sigma' a_f)^2 m_Z}{64\pi^2 \Gamma_Z} \left| \sum_\lambda \mathcal{M}_\sigma^\lambda(\theta) d_\lambda^{\sigma'}(\hat{\theta}, \hat{\varphi}) \right|^2, \quad (5)$$

where use has been made of the limit $\Gamma_Z \ll m_Z$, $\hat{\theta}$ and $\hat{\varphi}$ are the fermion angles in the center-of-mass frame of the decaying Z , and

$$\begin{aligned}d_{(\lambda=0)}^{\sigma'}(\hat{\theta}, \hat{\varphi}) &= \sigma' \sin \hat{\theta} \\ d_{(\lambda=\pm 1)}^{\sigma'}(\hat{\theta}, \hat{\varphi}) &= \frac{1}{\sqrt{2}} (1 \pm \sigma' \cos \hat{\theta}) e^{\pm i\hat{\varphi}}.\end{aligned}\quad (6)$$

We expand the squared matrix elements and define asymmetries:

$$\begin{aligned}\left| \sum_\lambda \mathcal{M}_\sigma^\lambda(\theta) d_\lambda^{\sigma'}(\hat{\theta}, \hat{\varphi}) \right|^2 &= \mathcal{F}_1(1 + \cos^2 \hat{\theta}) + \mathcal{F}_2(1 - 3\cos^2 \hat{\theta}) + \mathcal{F}_3 \cos \hat{\theta} + \\ &\quad \mathcal{F}_4 \sin \hat{\theta} \cos \hat{\varphi} + \mathcal{F}_5 \sin(2\hat{\theta}) \cos \hat{\varphi} + \mathcal{F}_6 \sin^2 \hat{\theta} \cos(2\hat{\varphi}) + \\ &\quad \mathcal{F}_7 \sin \hat{\theta} \sin \hat{\varphi} + \mathcal{F}_8 \sin(2\hat{\theta}) \sin \hat{\varphi} + \mathcal{F}_9 \sin^2 \hat{\theta} \sin(2\hat{\varphi}).\end{aligned}\quad (7)$$

$$A_i = \int_{-1}^1 d\cos\theta \mathcal{F}_i / \int_{-1}^1 d\cos\theta \mathcal{F}_1, \quad (8)$$

$$A'_i = \left(\int_0^1 - \int_{-1}^0 \right) d\cos\theta \mathcal{F}_i / \int_{-1}^1 d\cos\theta \mathcal{F}_1. \quad (9)$$

Table 1: CP and CPT^7 properties of squared matrix elements, and the corresponding observable asymmetries. CP and CPT conservation is indicated with a +, nonconservation is indicated with a -. The circles indicate that observation of the asymmetry requires identification of the charge of the final fermion f . The triangles indicate that the asymmetry may be suppressed without corresponding polarization measurements.

Matrix Elements	Properties CP CPT		Observables	beam Pol.	f Pol.	f charge
$ \widehat{M}_\sigma^+ ^2 + \widehat{M}_\sigma^- ^2 + \widehat{M}_\sigma^0 ^2$	+	+	σ_{tot}	-	-	-
$ \widehat{M}_\sigma^0 ^2$	+	+	A_2	-	-	-
$ \widehat{M}_\sigma^+ ^2 - \widehat{M}_\sigma^- ^2$	-	-	A'_1	$\triangle$	-	-
			A_3	-	$\triangle$	$\bigcirc$
$\text{Re}[(\widehat{M}_\sigma^+ - \widehat{M}_\sigma^-)(\widehat{M}_\sigma^0)^*]$	+	+	A_4	$\triangle$	$\triangle$	$\bigcirc$
			A'_5	-	-	-
$\text{Re}[(\widehat{M}_\sigma^+ - \widehat{M}_\sigma^-)(\widehat{M}_\sigma^0)^*]$	-	-	A'_4	-	$\triangle$	$\bigcirc$
			A_5	$\triangle$	-	-
$\text{Re}[(\widehat{M}_\sigma^+)(\widehat{M}_\sigma^-)^*]$	+	+	A_6	-	-	-
$\text{Im}[(\widehat{M}_\sigma^+ - \widehat{M}_\sigma^-)(\widehat{M}_\sigma^0)^*]$	-	+	A_7	$\triangle$	$\triangle$	$\bigcirc$
			A'_8	-	-	-
$\text{Im}[(\widehat{M}_\sigma^+ + \widehat{M}_\sigma^-)(\widehat{M}_\sigma^0)^*]$	+	-	A'_7	-	$\triangle$	$\bigcirc$
			A_8	$\triangle$	-	-
$\text{Im}[(\widehat{M}_\sigma^+)(\widehat{M}_\sigma^-)^*]$	-	+	A_9	-	-	-

It is remarkable that for each combination of matrix elements, one asymmetry exists which requires measurement neither of the final fermion spin nor of its charge, and hence all the visible Z decay modes can be used for these measurements. We present in Table 1 the nine combinations of matrix elements which appear. The asymmetries, averaged over electron and summed over fermion spins, are shown in Fig. 1 as deviations from the Standard Model value $a_Z = g_Z m_Z/2$. In Fig. 1a, the added couplings here are b_Z and c_Z ; the Higgs is a pure scalar in this plot. Taking instead a pure pseudoscalar, that is, taking only $\tilde{b}_Z$ to be nonzero, we would find a very different result. Then $A_2 = A_4 = A'_5 = 0$, and $A_6 \simeq -0.42$. Fig. 1b shows a few of the CP-violating asymmetries. Fermion angular correlations in the process $e^+e^- \rightarrow f\bar{f}\phi$ are useful in obtaining detailed information on the CP nature of a spinless neutral particle ϕ .

3. Acknowledgements

The authors wish to thank Peter Zerwas for discussions that prompted us to investigate the problem of measuring the 'Higgs' boson couplings. One of us (MLS) would also like to thank the NSF's Summer Institute in Japan program.




Figure 1: Asymmetries of Eqs. (8) and (9) for $m_H = 60$ GeV and $\sqrt{s} = 200$ GeV. The solid lines indicate the effect of b_Z , the dashed lines give the effect of c_Z . (a) A_2 – A_6 exist in the Standard Model. (b) A_7 – A_9 indicate CP violation.

This work was funded in part by the University of Wisconsin Research Committee with funds granted by the Wisconsin Alumni Research Foundation, and in part by the U.S. Department of Energy under Contract No. DE-AC02-76ER00881.

4. References

1. Proceedings of the Workshop "Physics and Experiments with Linear Colliders", edited by R. Orava, P. Eerola and M. Nordberg (World Scientific, 1992); Proceedings of the Workshop " e^+e^- Colliders at 500 GeV: The Physics Potential", DESY 92-123, edited by P. M. Zerwas; JLC-I, KEK Report 92-16.
2. J. F. Gunion, H. E. Haber, G. L. Kane, and S. Dawson, *The Higgs Hunter's Guide*, Addison-Wesley, 1990.
3. J. Fleischer and F. Jegerlehner, Nucl. Phys. **B216**, 469 (1983); B. Kniehl, Z. Phys. **C55**, 605 (1992); A. Denner, J. Küblbeck, R. Mertig, and M. Böhm, Z. Phys. **C56**, 261 (1992); S. Dawson and H.E. Haber, Phys. Rev. **D44**, 53 (1991).
4. A. Ali and M.A.B. Beg, Phys. Lett. **103B**, 376 (1981).
5. S. Weinberg, Phys. Rev. **D7**, 1068 (1973); Phys. Rev. **D42**, 860 (1990); B. Grzadkowski and J. F. Gunion, Phys. Lett. **B294**, 361 (1992).
6. K. Hagiwara and D. Zeppenfeld, Nucl. Phys. **B274**, 1 (1986).
7. K. Hagiwara, R. D. Peccei, D. Zeppenfeld and K. Hikasa, Nucl. Phys. **B282**, 253 (1987).